

20 Tips

to Help Prevent Medical Errors In Children

What Can You Do?

Be Involved in Your Child's Health Care

1. The single most important way you can help to prevent errors is to be an active member of your child's health care team.

Medicines

2. Make sure that all of your child's doctors know about everything your child is taking and his or her weight.

At least once a year, bring all of your child's medicines and supplements with you to the doctor. "Brown bagging" your child's medicines can help you and your doctor talk about them and find out if there are any problems.

3. Make sure your child's doctor knows about any allergies and how your child reacts to medicines.

This can help you avoid getting a medicine that can harm your child.

4. When your child's doctor writes you a prescription, make sure you can read it.

If you can't read the doctor's handwriting, your pharmacist might not be able to either. Ask the doctor to use block letters to print the name of the drug.

5. When you pick up your child's medicine from the pharmacy, ask: Is this the medicine that my child's doctor prescribed?

6. Ask for information about your child's medicines in terms you can understand- both when the medicines are prescribed and when you receive them at the hospital or pharmacy.

What is the name of the medicine? What is the medicine for? Is the dose of this medicine appropriate for my child based on his or her weight? How often is my child supposed to take it, and for how long? What side effects are likely? What do I do if they occur? Is this medicine safe for my child to take with other medicines or dietary supplements? What food, drink, or activities should my child avoid while taking this medicine? Is the dose of this medicine appropriate for my child based on his or her weight? When should I see an improvement?

7. If you have any questions about the directions on your child's medicine labels, ask.

Medicine labels can be hard to understand. For example, ask if "four doses daily" means taking a dose every 6 hours around the clock or just during regular waking hours.

8. Ask your pharmacist for the best device to measure your child's liquid medicine. Also, ask questions if you're not sure how to use the device.

Research shows that many people do not understand the right way to measure liquid medicines. For example, many use household teaspoons, which often do not hold a true teaspoon of liquid. Special devices, like marked oral syringes, help people to measure the right dose. Being told how to use the devices helps even more.

9. Ask for written information about the side effects your child's medicine could cause.

If you know what might happen, you will be better prepared if it does-or, if something unexpected happens instead. That way, you can report the problem right away and get help before it gets worse.

Hospital Stays

10. If you have a choice, choose a hospital at which many children have the procedure or surgery your child needs.

Research shows that patients tend to have better results when they are treated in hospitals that have a great deal of experience with their condition. Find out how many of the procedures have been performed at the hospital. While your child is in the hospital, make sure he or she is always wearing an identification bracelet.

11. If your child is in the hospital, ask all health care workers who have direct contact with your child whether they have washed their hands.

Handwashing is an important way to prevent the spread of infections in hospitals. Yet, it is not done regularly or thoroughly enough.

12. When your child is being discharged from the hospital, ask his or her doctor to explain the treatment plan you will use at home.

This includes learning about your child's medicines and finding out when he or she can get back to regular activities.

Surgery

13. If your child is having surgery, make sure that you, your child's doctor, and the surgeon all agree and are clear on exactly what will be done.

Doing surgery at the wrong site (for example, operating on the left knee instead of the right) is rare-but even once is too often. The good news is that wrong-site surgery is 100 percent preventable. The American Academy of Orthopaedic Surgeons urges its members to sign their initials directly on the site to be operated on before the surgery.

Other Steps You Can Take

14. Speak up if you have questions or concerns.

You have a right to question anyone who is involved with your child's care.

15. Make sure that you know who (such as your child's pediatrician) is in charge of his or her care.

16. Make sure that all health professionals involved in your child's care have important health information about him or her.

Do not assume that everyone knows everything they need to. Don't be afraid to speak up.

17. Ask a family member or friend to be there with you and to be your advocate.

Choose someone who can help get things done and speak up for you if you can't.

18. Ask why each test or procedure is being done.

It is a good idea to find out why a test or treatment is needed and how it can help. Your child could be better off without it.

19. If your child has a test, ask when the results will be.

If you don't, hear from the doctor or the lab, call to ask about the test results.

20. Learn about your child's condition and treatments by asking the doctor and nurse and by using other reliable sources.

Ask your child's doctor if his or her treatment is based on the latest scientific evidence.

Distributed by:

PULSE of America Inc.
Persons United Limiting Substandards and Errors
in Healthcare

1-800-96PULSE - www.PULSEAmerica.org

The 20 Tips To Help Prevent Medical Errors In Children are provided by the Agency for Healthcare Research and Quality.